

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/334082693>

Growth and survival of silver pompano *Trachinotus blochii* (Lacepede, 1801) at different salinities in inland saline ground water

Article in *The Indian journal of animal sciences* · May 2019

DOI: 10.56093/ijans.v89i5.90033

CITATIONS

9

READS

1,870

6 authors, including:



Madhuri Pathak

Central Institute of Fisheries Education

7 PUBLICATIONS 20 CITATIONS

[SEE PROFILE](#)



Wazir Singh Lakra

Central Institute of Fisheries Education

571 PUBLICATIONS 6,002 CITATIONS

[SEE PROFILE](#)



Appidi Krishna Reddy

Central Institute of Fisheries Education

74 PUBLICATIONS 669 CITATIONS

[SEE PROFILE](#)



Narinder Kumar Chadha

Central Institute of Fisheries Education

186 PUBLICATIONS 1,143 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Hormonal manipulation on testicular maturation and spermiation of *Clarias magur* (Hamilton 1822) [View project](#)



Enhancing the Utilisation of Nonfood Fish and Seaweed through Technology Refinement and Upscaling [View project](#)



Growth and survival of silver pompano *Trachinotus blochii* (Lacepede, 1801) at different salinities in inland saline ground water

MADHURI S PATHAK¹, WAZIR S LAKRA², APPIDI K REDDY³, N K CHADHA⁴, V K TIWARI⁵ and P P SRIVASTAVA⁶

ICAR-Central Institute of Fisheries Education, Mumbai, Maharashtra 400 061 India

Received: 19 September 2018; Accepted: 31 October 2018

ABSTRACT

Inland saline ground water (ISGW) has attracted great interest for commercial aquaculture of marine species in Haryana and Punjab. The silver pompano (*Trachinotus blochii*), a candidate for mariculture, was investigated for its potential to survive in inland saline water at ICAR-CIFE, Regional Centre, Rohtak, Haryana during 2015–16. The experimental set up comprised ISGW without potassium amendment, ISGW with potassium amendment and artificial sea water of 5 ppt, 10 ppt and 15 ppt salinity. In ISGW without potassium amendment, 100% mortality was observed at 12 h (5 ppt), 24 h (10 ppt) and 48 h (15 ppt). In the second experiment, three treatments (ISW5, ISW10 and ISW15) of ISGW with potassium amendment equivalent to sea water were prepared. A 100% survival was observed at all salinities in ISGW with potassium amendment, similar to the artificial seawater at the end of 90 days. Significantly higher weight gain, specific growth rate (SGR) and average daily growth (ADG) were observed at ISW10 and ISW15 than at ISW5. The plasma osmolality of silver pompano in ISW10 and ISW15 indicated that silver pompano regulates isosmotic condition to maintain the homeostasis of body. The present results indicate that silver pompano can be reared at 10 ppt to 15 ppt salinity with optimum growth and survival in potassium amended ISGW.

Key words: Inland saline aquaculture, Silver pompano, *Trachinotus blochii*

The salinization of agricultural land is a global concern for both developing as well as the developed nations. Saline ground water is not suitable for traditional agriculture (Joshi and Tyagi 1994, Forseberg *et al.* 1996, Allan *et al.* 2001) hence its use for aquaculture of marine water species can be a potential alternative with significant economic gains especially in India because of availability of huge volume of saline ground water resources especially in north-western parts of the country. In India, the salt affected lands are spread over an area of 8.62 m ha with 0.620 m ha in Haryana (Lakra *et al.* 2014). Ionic concentration of inland saline ground water (ISGW) is different from natural sea water in terms of its specific ions, which varies from location to location (Forsberg and Neill 1997). ISGW on the Indian sub-continent usually contains comparatively higher concentration of calcium than magnesium and potassium (Jain *et al.* 2002). The difference in the ionic concentration of ISGW affects the survival and growth of fish and shrimp

species (Fielder *et al.* 2001), as potassium plays very important role in osmo- and iono-regulation and acid-base balance in fishes (Marshall and Bryson 1998, Evans *et al.* 2005). India has achieved remarkable success in the farming of *Penaeus monodon* (Lakra *et al.* 2013a, Purushothaman *et al.* 2014) and *Litopenaeus vannamei* (Lakra *et al.* 2013b) at Rohtak, Haryana. Successful experiments were also conducted in the past to evaluate the effects of ionic deficiency and feasibility of marine fish culture in ISGW (Dwivedi and Lingaraju 1986, Jain *et al.* 2002, Raizada *et al.* 2005, Jain *et al.* 2006).

The silver pompano (*Trachinotus blochii*) with its fast growth rate is an important aquaculture species and has a very high market demand due to its good meat quality (Gopakumar *et al.* 2011). In India, ICAR-Central Marine Fisheries Research Institute has successfully carried out broodstock development, induced breeding and larval production of this species in 2011 (Gopakumar *et al.* 2011). The silver pompano is able to acclimatize and grow well even at a low salinity of about 8 ppt (Gopakumar *et al.* 2011). However, the potential of silver pompano culture in inland saline ground water is not known. Therefore, a study was conducted to evaluate the survival, growth and osmolality of the juveniles reared in potassium amended ISGW of varying salinities (5 ppt, 10 ppt and 15 ppt) compared to reconstituted seawater of similar salinities.

Present address: ¹Scientist (madhuri@cife.edu.in), ⁴Principal Scientist and Head (nkchadha@cife.edu.in), ⁵Principal Scientist (vktiwari@cife.edu.in), Division of Aquaculture; ⁶Principal Scientist (ppsrivastava63@gmail.com), Division of Fish Nutrition, Biochemistry and Physiology; ²Former Director (wslakra@gmail.com), ³Emeritus Scientist (akreddy.aqua@gmail.com).

MATERIALS AND METHODS

Experimental design: The experiment was conducted at wet lab complex of CIFE, Rohtak Centre, Haryana, India. Silver pompano (*Trachinotus blochii*) juveniles with an average length and weight of 6 cm and 4.5 g respectively, were procured from Mandapam regional center of Central Marine Fisheries Research Institute. Two sets of experiments were carried out to see the feasibility of silver pompano rearing in inland saline ground water (ISGW). In the first experiment, pompano juveniles were reared in unamended ISGW (RSW) at 5 ppt, 10 ppt and 15 ppt salinity to evaluate the effect of ISGW without any potassium amendment on its survival. Second experiment was conducted to determine the effect of potassium amendment in ISGW on the survival and growth of silver pompano juveniles. To make potassium amendment in ISGW, experimental media were prepared by supplementing commercial grade Muriate of Potash (KCl) with 100% equivalence potassium concentration of seawater (35 ppt) diluted to 5 ppt, 10 ppt and 15 ppt salinity, i.e. ISW5, ISW10 and ISW15 respectively. Reconstituted artificial sea water of 5 ppt, 10 ppt and 15 ppt salinities were prepared as reference media, i.e. ASW5, ASW10 and ASW15 respectively. Ionic profile of treatment and reference medium is given in Table 1. The experimental design used was a completely randomized design (CRD) wherein the reference and treatments were studied in triplicate. A total of 27 FRP tanks of 300 litre capacity were used in the experiment. The tanks were filled with 200 litres of treatment water and fitted with aeration stones to provide continuous aeration. A total of 15 juveniles (~4.5 g) pre-acclimatized to their respective salinities were stocked in each tank. The experiment was carried out for 90 days. Feeding was done using commercial Growel feed crumbles of size 1.8 mm three times a day. Water exchange was carried out once in a day during evening hours wherein 50% of water was exchanged. All the experimental media were prepared 15 days prior to the commencement of the trial. The salinity of the water was tested using a handheld refractometer (Atago, Tokyo, Japan) and was counter checked with titration method (APHA 2005). The Na²⁺ and K⁺ levels were analyzed using flame photometry

(Microprocessor flame photometer, Model 1382, ESICO, Haryana, India). The total hardness, Ca²⁺ and Mg²⁺ concentrations were analyzed using ethylene diamine tetra-acetic acid (EDTA) titration (APHA 2005).

Survival and growth performance: For the first experiment, survival of juveniles in unamended inland saline ground water (RSW) was recorded at time intervals of 6, 8, 12, 16, 20, 24, 48 and 72 h. During the second experiment, survival and growth were monitored for the juveniles reared in amended ISGW by counting and weighing the fishes every 7th day till the commencement of the experiment.

Percentage survival (PS), weight gain (WG), specific growth rate (SGR) were calculated as per the following formulas:

$$PS (\%) = \frac{\text{Number of survivors at the end of the experiment}}{\text{Number stocked initially}} \times 100$$

$$WG = \frac{W_2 - W_1}{W_1}$$

$$SGR = (\% \text{ body weight/day}) = \frac{\ln(W_2 - W_1)}{\Delta t} \times 100$$

$$ADG = \frac{W_2 \times W_1}{\Delta t}$$

where W₁, initial wet weight of fish at stocking; W₂, final wet weight of fish; t, grow out period.

Plasma and water osmolality: Fishes were anaesthetized using clove oil and blood was extracted from the caudal arch. Blood samples were centrifuged at 3,000 rpm for 3 min and the plasma thus obtained was used for osmolality measurement. Water samples from all the treatments were collected to measure osmolality. The water and plasma osmolality (mOsm/kg) were measured using a cryoscopic osmometer (Osmomat® 030, Gonotec GmbH, Germany). The osmoregulatory capacity (OC) of the juveniles in different treatments was calculated as the difference between the mean osmolality of the fish plasma and mean osmolality of their corresponding rearing media (Greenwell *et al.* 2003).

Statistical analysis: Statistical analysis were performed

Table 1. Ionic profile of treatment and reference medium used for 90 days trial (Mean±SE)

Treatment	Salinity (ppt)	Na ⁺ (mg/l)	K ⁺ (mg/l)	Mg ²⁺ (mg/l)	Ca ²⁺ (mg/l)	Na ⁺ /K ⁺ ratio	Mg ²⁺ /Ca ²⁺ ratio	K ⁺ (%)
RSW5	5.1±0.10	1393.67	4.5	306.67	156	309.70	1.97	8.4
RSW10	10.2±0.10	2703	8.1	541	314.33	333.70	1.72	7.5
RSW15	15.1±0.10	3712	11.5	767	413	322.78	1.86	7.17
ISW5	5.2±0.10	1394	55.33	396.67	158	25.19	2.51	103.42
ISW10	10.3±0.10	2765	109.43	789.33	320.33	25.27	2.46	102.27
ISW15	15.2±0.10	4066.33	161.83	1016.67	402.67	25.13	2.52	100.83
ASW5	5.1±0.10	1490	54.03	211	59.28	27.58	3.56	100.99
ASW10	10.3±0.10	3043.33	107.89	389.67	110.33	28.21	3.53	100.83
ASW15	15.0±0.20	4467	162.05	554	158.67	27.57	3.49	100.97

using SPSS, version 22. Treatment means were compared with one-way ANOVA and differences evaluated using Duncan's multiple range test. A significance level of $P < 0.05$ was used.

RESULTS AND DISCUSSION

During the experimental period of 90 days, all the water quality parameters did not differ among the treatments (Table 2). Mean values of all water quality parameters in treatment medium with potassium amendment (ISW) and reference medium reconstituted sea water (ASW) at 5 ppt, 10 ppt and 15 ppt were within the acceptable ranges for pompano culture (Jayakumar *et al.* 2014, Hamed *et al.* 2016). Except potassium concentration in ISGW, all the major ions were at par with sea water of equivalent salinity.

Inland saline ground water (ISGW) without any potassium amendment (RSW) was not suitable for the growth and survival of silver pompano (*T. blochii*), as the first experiment of feasibility study in RSW of all the experimental salinities resulted in 100% mortality. In 5 ppt RSW, juveniles became motionless in first 3 h, abnormal swimming was observed until 6 h and all the juveniles died before 12 h. At 10 ppt RSW, fishes exhibited normal swimming until 8 h. After 12 h juveniles lost the balance and complete mortality was observed in 24 h. However in 15 ppt RSW, survival was recorded for 48 h. During the current study, mortality of juveniles in RSW at different experimental salinities could not be only as a result of salinity change alone, as *T. Blochii* is a euryhaline fish (Kalidas *et al.* 2012). The difference in survival period in RSW could be attributed to osmotic stress caused by variations in ionic concentration of ISGW as compared with sea water of equivalent salinities. Several authors have also reported that, ion deficiency is a kind of stress which can have several effects on the fish including; changes in metabolic rate, health, behaviour, growth, survival and reproductive success (Brett 1958, Wedemeyer *et al.* 1976, Esch and Hazen 1980, Barton *et al.* 1990). It is also assumed that animals will grow best in low salinity water with ion ratios similar to that of diluted seawater (Boyd 2006). Potassium plays an important role in many physiological processes by maintaining the electrolyte and acid base balance (Wilson and El Naggar 1992). Deficient potassium

in inland saline water caused mortality in a number of cultured marine finfish species. Fielder *et al.* (2001) reported mortality of Australian snapper (*Pagrus auratus*) juveniles in 4 days when stocked in 19 ppt raw saline groundwater to the low concentration of potassium (5% K equivalence). Partridge and Creeper (2004) also stated mortality caused by hypokalemic muscle myopathy in barramundi, *Lates calcarifer*, reared in 45 ppt groundwater containing 25% of the potassium found in equivalent salinity seawater (25% K equivalence). Doroudi *et al.* (2006) observed the failure of juvenile mullet to survive at a low potassium concentration in inland saline groundwater. Saoud *et al.* (2012) reported that potassium-deficient low salinity well water is also not suitable for long-term culture of *Siganus rivulatus*. In case of crustaceans, it's a very well established fact that potassium ion deficiency in ISGW has caused total mortality in *P. monodon* (Raizada *et al.* 2003, Rahman *et al.* 2005) and *Litopenaeus vannamei* (Lakra *et al.* 2013, Pathak *et al.* 2013). This problem was solved with ionic supplementation of potassium for *P. monodon* (Raizada *et al.* 2003a, Rahman *et al.* 2005, Tantulo and Fotedar 2006) and other marine shrimps (Saoud *et al.* 2003, McNevin *et al.* 2004, Pragnell and Fotedar 2005, Bartholomew 2008, Pathak *et al.* 2013). However, initial experiments of rearing brackish water fishes in subsoil water indicated that culture of *Eetroplus suratensis* and *Mugil cephalus* is suitable without potassium supplementation (Dwivedi and Lingaraju 1986). Raizada *et al.* (2005) also reported that the milkfish *Chanos chanos* was also grown well in inland saline ground water of 18–25 ppt. Even in some freshwater fishes such as *Pangasius hypophthalmus*, Red tilapia, a potassium amendment does not have much influence on survival and growth as reported by Kumar *et al.* (2013) and Poonam *et al.* (2015) respectively. The experiment conducted by Ingram *et al.* (2002) achieved 97% survival of silver perch, *Bidyanus bidyanus*, in 10 ppt ground water with 22% K equivalence, highlights that tolerance to potassium deficiency is species specific. It was therefore necessary to amend potassium in ISGW for survival and growth of silver pompano.

In the second experiment, survival of juveniles in amended inland saline ground water was not affected by experimental salinities, which was close to 100% in all

Table 2. Water quality parameters of treatment and reference medium holding *Trachinotus blochii* (silver pompano) juveniles at different salinities during 90 days trial, expressed as Mean±SE

Water quality parameter	Treatment medium			Reference medium		
	ISW 10	ISW 15	ISW 15	ASW 5	ASW 10	ASW 15
Dissolved oxygen (mg/l)	7.10±0.21	6.83±0.33	6.77±0.20	7.00±0.12	7.03±0.07	7.10±0.21
Temperature (°C)	27.83±0.19	27.67±0.27	28.53±0.43	28.80±0.31	26.75±9.15	28.17±0.15
Free CO ₂ (mg/l)	0.85±0.20	0.91±0.23	0.64±0.17	0.78±0.21	0.85±0.31	0.91±0.21
pH	8.20±0.06	8.23±0.15	8.20±0.06	8.33±0.12	8.37±0.09	8.30±0.17
Alkalinity (mg/l)	195.33±9.33	218.00±4.36	242.33±4.48	182.33±9.91	228.00±6.81	249.67±19.10
Hardness (mg/l)	1342±10.12	2398.33±49.69	3239.67±72.13	1517.67±47.42	2351±72	3357.67±116
NH ₃ -N (mg/l)	0.03±0.00	0.06±0.01	0.08±0.01	0.07±0.01	0.05±0.01	0.04±0.01
NO ₂ -N (mg/l)	0.004±0.001	0.003±0.001	0.003±0.001	0.002±0.001	0.003±0.000	0.002±0.001

treatments ($P>0.05$) where potassium was amended at 100% equivalence to sea water of similar salinity. This was in agreement with the previous silver pompano research conducted by Kalidas *et al.* (2012) who reported minimum salinity required for survival of juveniles as 4 ppt. Sampaio *et al.* (2003) observed that pompano tolerate wide range of salinity, between 7 and 58 ppt on acute exposure of individuals acclimated to seawater (35 ppt), and on gradual exposure to diluted lower salinity sea water. The experiments conducted by Allen and Avault (1970) and Kumpf (1971) also reported that juvenile pompano were able to grow at a salinity of 5 ppt and were able to tolerate salinities as low as 2 ppt and as high as 45 ppt respectively. In the present study, the initial average body weight of silver pompano juveniles was not significantly different ($P<0.05$) at the commencement of the experiment. However, final mean individual weight, weight gain and SGR were significantly different towards the end of the trial (Table 3). Silver pompano juveniles exhibited significantly lower final mean body weight of juveniles in ISW 5 (14.31 g) and ASW 5 (15.59 g) than those of their counterparts in other treatments and reference media respectively. Highest final mean body weight was observed in ISW10 (36.12 g) followed by ISW 15 (35.85 g), but did not differ significantly to each other ($P>0.05$), similar trend was observed for weight gain, SGR and ADG. The growth of silver pompano juveniles in 10 and 15 ppt ISW and ASW showed no significant difference though numerically higher values were obtained at 10 ppt ISW and 15 ASW. Highest growth of pompano in 15 ASW medium was observed similar to the trend reported by Kalidas *et al.* (2012), who recommended salinity range for silver pompano in low saline water as 15–25 ppt.

Growth is the net positive result from the energy provided by food ingestion and the metabolic expenditure (Jobling 1994) hence, the growth of cultured fish would be maximum when they are reared in salinity near the isosmotic condition as the osmoregulatory cost is proportional to the osmotic gradient existing between the fish body fluids and the external medium (Handeland *et al.* 1998). Brown (2007) evaluated growth of Southern flounder, *Paralichthys*

lethostigma in inland low saline water and reported that at lower salinities (2.7 ppt and 3.4 ppt), the growth was very low as less energy is available for growth of fish. Instead of expending energy on somatic growth, fishes are constantly adjusting to their external surrounding environments and maintaining the proper balance of salt solutions within their bodies (isosmotic) by osmoregulation (Lisboa *et al.* 2015). Osmolality of silver pompano at 5 ppt exhibited a stressful condition as plasma osmolality of silver pompano juveniles in ISW 5 and ASW 5 (158.67 mmol/kg and 197.33 mmol/kg respectively) was significantly different than their medium osmolality (251.67 mmol/kg and 246 mmol/kg) respectively after 90 days. The lower growth observed in silver pompano juvenile maintained at 5 ppt ISGW in the present study seems to be related to a higher energy cost associated with osmoregulation. Fishes reared in hypo-osmotic environments would show additional energy requirements for osmoregulation that could hamper growth when compared to the fishes kept at an isosmotic environment (Boeuf and Payan 2001, Tsuzuki *et al.* 2007, Herrera *et al.* 2009, Perez-Robles *et al.* 2012). Laiz-Carrion *et al.* (2005) also reported that gilthead sea bream exhibited osmoregulatory difficulties when reared at 4 ppt due to the lower plasma osmolality after 42 and 84 days and in European sea bass after 24 h of transfer to salinities less than 20 ppt (Venturini *et al.* 1992). Low salinity has been reported to cause difficulties in homeostasis in other euryhaline teleost species, such as the sheepshead minnow *Cyprinodon variegatus* (Nordlie, 1985) and flounder *Paralichthys orbignyanus* (Sampaio and Bianchini 2002). Osmoregulatory imbalance indicates the salinity stress (Franklin *et al.* 1992), and negative effects of stress on growth are well documented in teleosts (WendelaarBonga 1997). Plasma osmolality of silver pompano juveniles in ISW 10 and ISW 15 at the end of experimental period (332.2 and 413 mmol/kg respectively) indicated that silver pompano regulated osmotic balance with treatment medium salinities. Similar trend was observed in plasma osmolalities of juveniles stocked in ASW 10 and ASW 15 (352.6 and 412.6 mmol/kg respectively). Plasma osmolality level of a marine fish is

Table 3. Performance of *Trachinotus blochii* (silver pompano) juveniles at different salinities of treatment and reference medium during 90 days trial, expressed as Mean \pm SE (n=3)

Parameter	Treatment medium			Reference medium		
	ISW 5	ISW 10	ISW 15	ASW 5	ASW 10	ASW 15
Initial body weight (g)	1.48 \pm 0.02	1.46 \pm 0.04	1.53 \pm 0.06	1.49 \pm 0.06	1.51 \pm 0.07	1.51 \pm 0.04
Final body weight (g)	14.31 \pm 2.55 ^a	36.12 \pm 2.87 ^b	35.85 \pm 2.35 ^b	15.59 \pm 1.64 ^a	37.42 \pm 3.66 ^b	39.13 \pm 1.18 ^b
Weight gain (g)	12.83 \pm 2.54 ^a	34.65 \pm 2.83 ^b	34.32 \pm 2.31 ^b	14.10 \pm 1.71 ^a	35.91 \pm 3.71 ^b	37.62 \pm 1.22 ^b
SGR (%/day)	2.48 \pm 0.19 ^a	3.55 \pm 0.06 ^b	3.50 \pm 0.06 ^b	2.60 \pm 0.17 ^a	3.56 \pm 0.15 ^b	3.62 \pm 0.06 ^b
ADG (g/f/d)	0.143 \pm 0.03 ^a	0.383 \pm 0.03 ^b	0.380 \pm 0.02 ^b	0.16 \pm 0.02 ^a	0.40 \pm 0.04 ^b	0.42 \pm 0.01 ^b
Survival (%)	100	100	100	100	100	100
Water osmolality (mOsm/kg)	251.67 \pm 19.6 ^a	345 \pm 10.2 ^b	446.67 \pm 13.3 ^c	246 \pm 21.7 ^a	372.33 \pm 12.5 ^b	433.67 \pm 10.4 ^c
Plasma osmolality (mOsm/kg)	158.67 \pm 9.68 ^a	332.33 \pm 4.91 ^c	413 \pm 5.69 ^d	197.33 \pm 4.06 ^b	352.67 \pm 9.53 ^c	412.67 \pm 13.1 ^c

^{a,b,c}Values marked with superscript letters show significant differences in the growth parameters among three treatments compared to reference medium ($P<0.05$).

typically maintained in between 380–450 mmol/kg (Conte 1969, Johnson 1973, Bond 1991, Kirschner 1991) and in teleosts, the plasma isosmotic point generally corresponds to the water salinity of 12 ppt (Boeuf and Payan 2001, Tsuzuki *et al.* 2007, Herrera *et al.* 2009, Nordlie 2009). Lowest osmoregularity capacity was observed at ISW 10 and ASW 10 (12.67 and 19.66 mmol/kg respectively) followed by ASW 15 and ISW 15 (21 and 33.67 mmol/kg respectively). Sarwono (2005) also reported in juvenile sea bass that the differences between blood and culture media osmolality in 10 and 15 ppt salinity were minimum, exhibiting the iso-osmotic point at 10 and 15 ppt. These results were in agreement with the fact that optimum growth of fish can be achieved when a particular range of salinity have the isotonic condition as the isotonic salinity has an effect on metabolic rates (Morgan and Iwana 1991). This suggests that silver pompano might have maintained isosmotic conditions at 10 ppt and 15 ppt ISW, thereby energetic loss for osmoregulation is lower associated with minimum gradients between water and blood. This enhanced the growth of silver pompano juveniles in these two treatments than at 5 ppt ISW.

Inland saline ground water is deficient in potassium and unsuitable for rearing silver pompano juveniles without potassium amendment. It has been observed that after potassium amendment, the growth was lower at 5 ppt with high osmotic capacity. However, the culture of silver pompano (*T. Blochi*) is feasible at 10 to 15 ppt inland saline ground water provided potassium amendment is done at par with that of sea water of equivalent salinity. Further studies should be oriented on physiological changes in silver pompano at different levels of potassium amendment.

ACKNOWLEDGEMENTS

The authors are thankful to Director, ICAR-Central Institute of Fisheries Education (CIFE), Mumbai for providing facilities for carrying out this work. The first author acknowledges the Indian Council of Agricultural Research (ICAR), New Delhi for the scholarship during the study period.

REFERENCES

- Allan G L, Banens L B and Fielder S. 2001. *Developing commercial inland saline aquaculture in Australia: part 2. Resource Inventory and Assessment*. FRDC Project No. 98/335, NSW Fisheries Final Report Series, No. 31, NSW, Australia.
- Allen K O and Avault J W Jr. 1970. Effects of salinity and water quality on survival and growth of juvenile pompano (*Trachinotus carolinus*). Coastal Studies Bulletin No. 5, Louisiana State University, Baton Rouge, LA. pp 147–155.
- APHA/WWA/WEF. 2005. *Standard Methods for the Examination of Water and Wastewater*. 21st edn. American Public Health Association, Washington, DC, USA.
- Bartholomew W G. 2008. Stocking strategies for production of *Litopenaeus vannamei* (Boone) in amended freshwater in inland ponds. *Aquaculture Research* **39**:10–17.
- Barton B A, McLeay D J and Wedemeyer G A. 1990. Stress and acclimation, pp. 451–89. *Methods for Fish Biology*. (Eds) Moyle P B and Schreck C B. American Fisheries Society, Bethesda, Maryland, USA.
- Boeuf G and Payan P. 2001. How should salinity influence fish growth? *Comparative Biochemistry and Physiology, Part C* **130**: 411–23.
- Bond C E. 1991. *Biology of Fishes*. 2nd edn. Saunders College Publishing, New York, USA.
- Boyd C A. 2006. 'Investigations of water supply and water quality issues related to inland shrimp farming in western Alabama'. Doctoral Dissertation, Auburn University, Auburn, Alabama, USA.
- Brett J R. 1958. Investigation of fish-power problems, pp. 69–83. *Implications and Assessment of Environmental Stress*. (Eds) Larkin A and MacMillan H R. Lectures in Fisheries, University of British Columbia, Vancouver, Canada.
- Brown B J. 2007. 'Evaluation of three fish species for culture using low salinity groundwater in the black belt region of Alabama'. Master Thesis, Auburn University, Alabama.
- Conte F P. 1969. Salt secretion, pp. 241–292. *Fish Physiology*. (Eds) Hoar W S and Randall D J. Volume 1. Academic Press, New York, USA.
- Doroudi M S, Fielder D S and Allan G L. 2006. Combined effects of salinity and potassium concentration on juvenile mullet (*Argyrosomus japonicus*, Temminck and Schlegel) in inland saline groundwater. *Aquaculture Research* **37**: 1034–39.
- Dwivedi S N and Lingaraju G M. 1986. Strategy for prawn and fish culture in saline sub soil waters of semi-arid zone of Haryana. *Mahasagar* **19**: 97–102.
- Esch G W and Hazen T C. 1980. Stress and body condition in a population of largemouth bass: implications for red sore disease. *Transactions of the American Fisheries Society* **110**: 644–49.
- Evans D H, Piermarini P M and Choe K P. 2005. The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid–base regulation, and excretion of nitrogenous waste. *Physiological Reviews* **85**: 97–177.
- Fielder D S, Bardsley W J and Allan G L. 2001. Survival and growth of Australian snapper, *Pagrus auratus*, in saline groundwater from inland New South Wales, Australia. *Aquaculture* **201**: 73–90.
- Forsberg J A and Neill W H. 1997. Saline groundwater as an aquaculture medium: physiological studies on the red drum, *Sciaenops ocellatus*. *Environmental Biology of Fishes* **49**: 19–128.
- Forsberg J A, Dorsett P W and Neill W H. 1996. Survival and growth of red drum *Sciaenops ocellatus* in saline groundwater of West Texas, USA. *Journal of the World Aquaculture Society* **27**: 462–74.
- Franklin C E, Forster M E and Davison W. 1992. Plasma cortisol and osmoregulatory changes in sockeye salmon transferred to sea water: comparison between successful and unsuccessful adaptation. *Journal of Fish Biology* **41**: 113–22.
- Gopakumar G, Syda Rao G, Abdul Nazar A K, Jayakumar R, Tamilmani G, Kalidas C, Sakthivel M, Rameshkumar P, Hanumantha R, Murugan A, Premjothi R, Balamurugan V, Ramkumar B and Jayasingh M. 2011. Silver pompano: A potential species for mariculture in India – breeding and seed production of silver pompano (*Trachinotus blochii*). *Fishing Chimes* **31**(6): 58–60.
- Greenwell G M, Johanna S and Clayton L A. 2003. Osmoregulation in fish. Mechanisms and clinical implications. The veterinary clinics of North America. *Exotic Animal*

- Practice* 6: 169–89
- Hamed S S, Narriman S J and Bwathondi P O J. 2016. Effect of salinity levels on growth, feed utilization, body composition and digestive enzymes activities of juvenile silver pompano (*Trachinotus blochii*). *International Journal of Fisheries and Aquatic Studies* 4(6): 279–83.
- Handeland S O, Berge A, Bjornsson B Th and Stefansson S O. 1998. Effect of temperature and salinity on osmoregulation and growth of Atlantic salmon (*Salmo salar* L.) smolts in seawater. *Aquaculture* 168: 289–302.
- Herrera M, Vargas-Chacoff L, Hachero I, Ruiz-Jarabo I, Rodiles A, Navas J I and Mancera J M. 2009. Osmoregulatory changes in wedge sole (*Dicologlossa cuneata* Moreau, 1881) after acclimation to different environmental salinities. *Aquaculture Research* 40: 762–71.
- Ingram B A, McKinnon L J and Gooley G J. 2002. Growth and survival of selected aquatic animals in two saline groundwater evaporation basins: an Australian case study. *Aquaculture Research* 33: 425–36.
- Jain A K, Kumar G and Mukherjee S C. 2006. Survival and growth of early juveniles of barramundi *Lates calcarifer* (Bloch, 1790) in inland saline groundwater. *Journal of Biological Research* 5: 93–97.
- Jain A K, Raju K D and Arasu A R T. 2002. Saline water resources of Rajasthan and their suitability for brackish water aquaculture. *Fishing Chimes* 22: 7–13.
- Jayakumar R, Abdul Nazar A K and Gopakumar G. 2014. *Culture of silver pompano (Trachinotus blochii) in coastal aquaculture ponds*. CMFRI Manual Customized Training Book. pp. 191–94.
- Jobling M. 1994. *Fish Bioenergetics*. London, Chapman and Hall, 309 p.
- Johnson D W. 1973. Endocrine control of hydromineral balance in teleosts. *American Zoology* 13: 799–818.
- Joshi P K and Tyagi N K. 1994. Salt affected and waterlogged soil in India: a review, pp. 237–252. *Strategic Change in Indian Irrigation*. (Eds) Svendsen M and Gulati A. ICAR, New Delhi and IFPRI, Washington, DC, USA.
- Kalidas C, Sakthivel M, Tamilmani G, Ramesh Kumar P, Abdul Nazar A K, Jayakumar R, Balamurugan, Ramkumar, Prem Jothi and Gopakumar G. 2012. Survival and growth of juvenile silver pompano (*Trachinotus blochii*) (Lacepède, 1801) at different salinities in tropical conditions. *Indian Journal of Fisheries* 59(3): 95–98.
- Kirschner L E. 1991. Water and ions, pp 13–107. *Environmental and Metabolic Animal Physiology: Comparative Animal Physiology*. 4th edn. (Ed) Prosser C L. Wiley-Liss, New York, USA.
- Kumar A, Harikrishna V, Reddy A K, Chadha N K and Lakra W S. 2013. 'Studies on growth and survival of *Pangasiodon hypophthalmus* in inland saline water'. MFSc. Dissertation, Central Institute of Fisheries Education, Versova, Mumbai.
- Kumpf H E. 1971. 'Temperature-salinity tolerance of the Florida pompano (*Trachinotus carolinus* Linnaeus)'. PhD Dissertation, University of Miami, Miami, FL.
- Laiç-Carrion R, Guerrero P M, Fuentes J, Canario A V M, Martin de Rio M P and Mancera J M. 2005. Branchial osmoregulatory response to salinity in the gilthead sea bream, *Sparus auratus*. *Journal of Experimental Zoology* 303A: 563–76.
- Lakra W S, Reddy A K and Harikrishna V. 2014. Technology for commercial farming of Pacific white shrimp *Litopenaeus vannamei* in inland saline soils using ground saline water. CIFE Technical Bulletin-1, pp.1–28.
- Lakra W S, Reddy A K, Harikrishna V and Ayyapan S. 2013. Farming of tiger shrimp in low saline water at Rohtak. *Fishing Chimes* 32(9): 45.
- Lakra W S, Reddy A K, Harikrishna V and Ayyapan S. 2013b. Farming of pacific white shrimp *Litopenaeus vannamei* in inland saline water ponds at Lahli farm of CIFE, Rohtak centre. *Fishing Chimes* 32(9): 44–45.
- Lisboa V, Indianara F B, Sampaio L A and Bianchini A. 2015. Effect of salinity on survival, growth and biochemical parameters in juvenile Lebranch mullet *Mugiliza* (Perciformes: Mugilidae). *Neotropical Ichthyology* 13: 447–52.
- Marshall W S and Bryson S E. 1998. Transport mechanisms of seawater teleost chloride cells: an inclusive model of a multifunctional cell. *Comparative Biochemistry and Physiology* 119A: 97–106.
- McNevin A A, Boyd C E, Silapajam O and Silapajam K. 2004. Ionic supplementation of pond waters for inland culture of marine shrimp. *Journal of World Aquaculture Society* 35: 460–67.
- Morgan J D and Iwama G K. 1991. Effects of salinity on growth, metabolism, and ion regulation in juvenile rainbow and steelhead trout (*Oncorhynchus mykiss*) and fall chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* 48(11): 2083–94.
- Nordlie F G. 1985. Osmotic regulation in the sheepshead minnow *Cyprinodon variegatus* Lacepede. *Journal of Fish Biology* 26: 161–70.
- Nordlie F G. 2009. Environmental influences on regulation of blood plasma serum components in teleost fish: a review. *Reviews in Fish Biology and Fisheries* 19: 481–564.
- Partridge G J and Creeper J. 2004. Skeletal myopathy in juvenile barramundi, *Lates calcarifer* (Bloch), cultured in potassium-deficient saline groundwater. *Journal of Fish Diseases* 27: 523–30.
- Pathak M S, Reddy A K, Harikrishna V, Chadha N K and Chandra Prakash. 2013. 'Ionic manipulation of inland saline ground water for growth and survival of *Litopenaeus vannamei*' (Boone, 1931). MFSc. Dissertation, Central Institute of Fisheries Education, Versova, Mumbai.
- Perez-Robles J, Re A D, Giffard-Mena I and Diaz F. 2012. Interactive effects of salinity on oxygen consumption, ammonium excretion, osmoregulation and Na⁺/K⁺-ATPase expression in the bullseye puffer (*Sphoeroides annulatus*, Jenyns 1842). *Aquaculture Research* 43: 1372–83.
- Poonam R, Harikrishna V, Lakra W S, Reddy A K and Babitha R A M. 2013. 'Performance evaluation of red tilapia in inland saline water at various salinity levels with special reference to ionic manipulation'. MFSc. Dissertation, Central Institute of Fisheries Education, Versova, Mumbai.
- Prangnell D I and Fotedar R. 2005. The effect of potassium concentration in inland saline water on the growth and survival of the western king shrimp, *Penaeus latisulcatus* (Kishinouye, 1896). *Journal of Applied Aquaculture* 17: 19–34.
- Purushothaman C S, Raizada S, Sharma V K, Harikrishna V, Venugopal G, Agrahari R K, Rahaman M, Hasan J and Kumar A. 2014. Production of tiger shrimp (*Penaeus monodon*) in potassium supplemented inland saline sub-surface water. *Journal of Applied Aquaculture* 26(1): 84–93.
- Rahman S, Jain A K, Reddy A K, Girish K and Koyya D R. 2005. Ionic manipulation of inland saline groundwater for enhancing survival and growth of *Penaeus monodon* (Fabricius). *Aquaculture Research* 36: 1149–56.

- Raizada S, Chadha N K, Maheshwari U K, Hasan J, Ali M, Singh I J and Kumar S. 2003. *Role of active metal cations on the survival and growth of tiger shrimp (Penaeus monodon) in inland ground saline water*. Proceedings of 3rd Interaction Workshop, Fish Production Using Brackishwater in Arid Eco System. (Eds) Garg S K and Arasu A R T. pp. 208–11, December 17–18, CCSHAU, Hisar, India.
- Raizada S, Chadha N K, Ali M, Kumar A and Javed H. 2005. Length-weight relationship of milkfish, *Chanos chanos* (Forsk.) reared in inland saline ground water. *Indian Journal of Fisheries* **52**: 115–17.
- Sampaio L A and Bianchini A. 2002. Salinity effects on osmoregulation and growth of the euryhaline Flounder *Paralichthys orbignyanus*. *Journal of Experimental Marine Biology and Ecology* **269**: 187–96.
- Sampaio L A, Tesser M B and Burkert D. 2003. Acute salinity tolerance of juvenile pompano *Trachinotus marginatus* (Teleostei, Carangidae) under laboratory conditions. *Ciencia Rural* **33**: 757–61.
- Saoud I P, Davis D A and Rouse D B. 2003. Suitability studies of inland well waters for *Litopenaeus vannamei* culture. *Aquaculture* **217**: 373–83.
- Saoud I P, Mourad N, Kreydiyyeh S and Ghanawi J. 2012. Aquaculture of marine fish in inland low salinity well water: potassium is not the only limiting element. *Fisheries and Aquaculture Journal* **10**: 2150–3508.
- Sarwono and Hidayat A. 2005. Effect of salinity on osmoregulatory capacity, feed consumption, feed efficiency and growth of juvenile sea bass (*Lates calcarifer* Bloch). Graduate School, Kasetsart University.
- Tantulo U and Fotedar R. 2006. Comparison of growth, osmoregulatory capacity, ionic regulation and organosomatic indices of black tiger shrimp (*Penaeus monodon* Fabricius, 1798) juveniles reared in potassium fortified inland saline water and ocean water at different salinities. *Aquaculture* **258**(1–4): 594–605.
- Tsuzuki M Y, Sugai J K, Maciel J C, Francisco C J and Cerqueira V R. 2007. Survival, growth and digestive enzyme activity of juveniles of the fat snook (*Centropomus parallelus*) reared at different salinities. *Aquaculture* **271**: 319–25.
- Venturini G, Cataldi E, Marino G, Pucci P, Garibaldi L, Bronzi P and Cataudella S. 1992. Serum ions concentration and ATPase activity in gills, kidney and oesophagus of European sea bass (*Dicentrarchus labrax*, Pisces, Perciformes) during acclimation trials to fresh water. *Comparative Biochemistry and Physiology* **103A**: 451–54.
- Wedemeyer G A, Meyer F P and Smith L. 1976. *Environmental Stress and Fish Diseases*. TFH Publications, Neptune City, New Jersey, USA.
- WendelaarBonga S E. 1997. The stress response in fish. *Physiological Reviews* **77**: 591–625.
- Wilson R P and Naggar E I G. 1992. Potassium requirement of fingerling channel catfish *Ictalurus punctatus*. *Aquaculture* **108**: 169–75.